

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of: Kenneth J. Rothschild et al.

Serial No.:

10/719,523

Filed:

11/21/03

Group No.: 1636

Examiner:

Schlapkohl, W.

Entitled:

Methods For The Detection, Analysis And Isolation Of Nascent Proteins

INFORMATION DISCLOSURE STATEMENT TRANSMITTAL

Mail Stop Amendment Commissioner for Patents P.O. Box 1450 Alexandria, VA 22313-1450

CERTIFICATE OF MAILING UNDER 37 C.F.R. § 1.8(a)(1)(i)(A)

I hereby certify that this correspondence (along with any referred to as being attached or enclosed) is, on the date shown below, being deposited with the U.S. Postal Service with sufficient postage as first class mail in an envelope addressed to: Mail Stop Amendment, Commissioner for Patents, P.O. Box 1450, Alexandria, VA

Dated: August 21, 2006

By:

Christopher J/Collin

Sir or Madam:

Enclosed please find an Information Disclosure Statement and Form PTO-1449, including copies of the references contained thereon, for filing in the U.S. Patent and Trademark Office.

A check for \$180.00 is also enclosed pursuant to 37 C.F.R. § 1.17(p) for filing this Information Disclosure Statement after three months as set forth in 37 C.F.R. § 1.97(c).

The Commissioner is hereby authorized to charge any additional fee or credit overpayment to our Deposit Account No. 08-1290. An originally executed duplicate of this transmittal is enclosed for this purpose.

Dated: August 21, 2006

Peter G. Carroll

Registration No. 32,837

MEDLEN & CARROLL, LLP 101 Howard Street, Suite 350 San Francisco, California 94105 617/984.0616

Group No.: 1636

Schlapkohl, W.



Application of: Kenneth J. Rothschild et al.

Serial No.: 10/719,523

Examiner: Filed: 11/21/03 Methods For The Detection, Analysis And Entitled:

Isolation Of Nascent Proteins

INFORMATION DISCLOSURE STATEMENT

Mail Stop Amendment Commissioner for Patents P.O. Box 1450 Alexandria, VA. 22313-1450

CERTIFICATE OF MAILING UNDER 37 C.F.R. § 1.8(a)(1)(i)(A)

I hereby certify that this correspondence (along with any referred to as being attached or enclosed) is, on the date shown below, being deposited with the U.S. Postal Service with sufficient postage as first class mail in an envelope addressed to: Mail Stop Amendment, Commissioner for Patents, P.O. Box 1450, Alexandria, VA. 2231

Dated: August 21, 2006

Sir or Madam:

The citations listed below, copies attached, may be material to the examination of the above-identified application, and are therefore submitted in compliance with the duty of disclosure defined in 37 C.F.R. §§ 1.56 and 1.97. The Examiner is requested to make these citations of official record in this application.

The following printed publications are referred to in the body of the specification:

- U.S. Pat. No. 4,683,195 to Mullis et al.;
- U.S. Pat. No. 4,774,339 to Haugland et al.;
- U.S. Pat. No. 5,069,769 to Fujimiya et al.;
- U.S. Pat. No. 5,091,328 to Miller;
- U.S. Pat. No. 5,137,609 to Manian et al.;
- U.S. Pat. No. 5,187,288 to Kang et al.;
- U.S. Pat. No. 5,190,632 to Fujimiya et al.;
- U.S. Pat. No. 5,248,782 to Haugland et al.;
- U.S. Pat. No. 5,274,113 to Kang et al.;
- U.S. Pat. No. 5,433,896 to Kang et al.;
- U.S. Pat. No. 5,451,663 to Kang et al.;
- U.S. Pat. No. 5,643,722 to Rothschild et al.;

08/23/2006 SFELEKE1 00000011 10719523

- U.S. Pat. No. 5,654,150 to King et al.;
- U.S. Pat. No. 5,783,397 to Hughes *et al.*;
- PCT WO90/05785 to Schultz;
- Allen et al., Gel Electrophoresis and Isoelectric Focusing of Proteins, Walter de Gruyter, New York 1984, pp.17-62;
- Antibodies: A Laboratory Manual (E. Harlow and D. Lane, editors, Cold Spring Harbor Laboratory Press, 1988) pp.53,72-73;
- Bain et al., "Site-Specific Incorporation of Nonnatural Residues during In Vitro Protein Biosynthesis with Semisynthetic Aminoacyl-tRNAs," Biochemistry 30:5411-21 (1991);
- Bruce and Uhlenbeck, "Specific Interaction of Anticodon Loop Residues with Yeast Phenylalanyl-tRNA Synthetase," *Biochemistry* 21:3921-3926 (1982);
- Current Protocols in Molecular Biology (F.M. Ausubel et al. editors, Wiley Interscience, 1993), 10-16,10-77;
- Da Poian, A. T., et al., "Kinetics of intracellular viral disassembly and processing probed by Bodily fluorescence dequenching," *J Virol Methods* 70(1), 45-58 (1998);
- Doty et al., "Strand Separation and Specific Recombination in Deoxyribonucleic Acids: Physical Chemicals Studies," Proc. Natl. Acad. Sci. USA 46:461 (1960);
- DiCesare et al., "A High-Sensitivity Electrochemiluminescence-Based Detection System for Automated PCR Product Quantitation," *BioTechniques* 15:152-59 (1993);
- Felgner *et al.*, "Lipofection: A highly efficient, lipid-mediated DNA-transfection procedure," *Proc. Natl. Acad. Sci. USA* 84:7413-17 (1987);
- Happ *et al.*, "New Approach to the Synthesis of 2'(3')-*O*-Aminoacyl Oligoribonucleotides," *J. Org. Chem.* 52:5387-91 (1987);
- Heckler et al., "Preparation of 2'(3')-O-Acyl-pCpA Derivatives as Substrates for T4 RNA Ligase-Mediated "Chemical Aminoacylation"," Tetrahedron 40:87-94 (1984);
- Heckler et al., "T4 RNA Ligase Mediated Preparation of Novel "Chemically Misacylated" tRNA Phes," Biochemistry 23:1468-73 (1984);
- Hemmila, I.A., <u>Chemical Analysis</u> "Applications of Fluorescence in Immunoassays", (Wiley&Sons 1991) pp.138-159;
- Hudson, "Methodological Implications of Simultaneous Solid-Phases Peptide Synthesis. 1. Comparison of Different Coupling Procedures," J. Org. Chem. 53:617-624 (1988);

- Krieg et al., "Photocrosslinking of the signal sequence of nascent preprolactin to the 54-kilodalton polypeptide of the signal recognition particle," *Proc. Natl. Acad. Sci. USA* 83:8604-08 (1986);
- Keller, R. C., et al., "Characterization of the Resonance Energy Transfer Couple Coumarin-Bodily and its Possible Applications in Protein-Lipid Research," Biochem Biophys Res Commun 207(2), 508-14 (1995);
- Kim, D., and Choi, C., "A Semicontinuous Prokaryotic Coupled Transcription/Translation System Using a Dialysis Membrane," *Biotechnol Prog* 12, 645-649 (1996);
- Kopp et al., "Chemical Amplification: Continuous Flow PCR on a Chip,"
 Science 280:1046 (1998);
- Kozak, "Point Mutations Define a Sequence Flanking the AUG Initiator Codon that Modulates Translation by Eukaryotic Ribosomes," Cell 44:283-292 (1986);
- Kudlicki, W.et al., "Chaperone-dependent Folding and Activation of Ribosomebound Nascent Rhodanese," J Mol Biol 244(3), 319-31 (1994);
- Laemmli, U. K., "Cleavage of Structural Proteins during the Assembly of the Head of Bacteriophage T4," *Nature* 227:680-685 (1970);
- Marmur and Lane, "Strand Separation and Specific Recombination in Deoxyribonucleic Acids: Biological Studies," *Proc. Natl. Acad. Sci. USA* 46:453-461 (1960);
- Molecular Cell Biology (J. Darnell et al. editors, Scientific American Books, N.Y., N.Y. 1991) pp.119-132;
- Neu and Heppel, "Nucleotide Sequence Analysis of Polyribonucleotide by Means of Periodate Oxidation Followed by Cleavage with an Amine," *J. Biol. Chem.* 239:2927-34 (1964);
- Noren *et al.*, "A General Method for Site-Specific Incorporation of Unnatural Amino Acids into Proteins," *Science* 244:182-188 (1989);
- Odom, O. W, et al., "In vitro engineering using acyl-derivatized tRNA," In Protein synthesis: Methods and Protocols, PP.93-103, (Humana Press, Totowa, NJ.);
- Patchornik et al., "Photosensitive Protecting Groups," J. Am. Chem. Soc. 92:6333-35 (1970);
- Pavlopoulos, et al., "Laser action from a tetramethylpyrromethene-BF.sub.2 complex," APP. OPTICS 27:4998-4999 (1988);

- Pfahler et al., Sensors and Actuators, A21-A23, pp. 431-434 (1990)¹;
- Pillai, "Photoremovable Protecting Groups in Organic Synthesis," *Synthesis* 1-26 (1980);
- Powell et al., "Molecular Diagnosis of Familial Adenomatous Polyposis," N. Engl. J. Med. 329:1982-87 (1993);
- Pratt, "Coupled Transcription-Translation in Prokaryotic Cell-Free System," (*Transcription and Translation*, B.D. Hames and S.J. Higgins, Editors, p. 179-209, IRL Press, Oxford, 1984);
- Promega Technical Bulletin No. 182; tRNA^{nscend}TM: Non-radioactive Translation Detection System, Sept. 1993;
- Reis, R. C., et al., "A novel methodology for the investigation of intracellular proteolytic processing in intact cells," Eur J Cell Biol 75(2), 192-7 (1998);
- Rowan and Bodmer, "Introduction of a myc Reporter Taq to Improve the Quality of Mutation Detection Using the Protein Truncation Test," Human Mutation 9:172-176 (1997);
- Sampson and Uhlenbeck, "Biochemical and physical characterization of an unmodified yeast phenylalanine transfer RNA transcribed in vitro," Proc. Natl. Acad. Sci. USA 85:1033-37 (1988);
- Seong and RajBhandary, "Escherichia coli formylmethione tRNA: Mutations in GGG sequence conserved in anticodon stem of initiator tRNAs affect initiation of protein synthesis and conformation of anticodon loop," Proc. Natl. Acad. Sci. USA 84:334-338 (1987);
- Spirin *et al.*, "A Continuous Cell-Free Translation System Capable of Producing Polypeptides in High Yield," *Sci.* 242:1162-64 (1988);
- Stephen, "High-Resolution Preparative SDS-Polyacrylamide Gel Electrophoresis: Fluorescent Visualization and Electrophoretic Elution-Concentration of Protein Bands," *Anal. Biochem.* 65:369-79 (1975);
- Treibs & Kreuzer, "Difluorboryl-komplexe von di- und tripyrrylmethenen,"
 Liebigs Ann. Chem. 718:208-223 (1968);
- Turcatti et al., "Probing the Structure and Function of the Tachykinin Neurokinin-2 Receptor through Biosynthetic Incorporation of Fluorescent Amino Acids at Specific Sites," J Biol Chem 271(33), 19991-8 (1996);

We have been unable to obtain this reference, but if the examiner requests a copy we will again seek to obtain it.

- Van Lintel et al., Sensors and Actuators 15:153-167 (1988)²;
- Varshney U. and RajBhandary UL, "Initiation of protein synthesis from a termination codon," Proc Natl Acad Sci U S A 87(4):1586-90 (1990);
- Varshney et al., "Direct Analysis of Aminoacylation Levels of tRNAa in Vivo," J. Biol. Chem. 266: 24712-24718 (1991);
- Yao S *et al.*, "SDS capillary gel electrophoresis of proteins in microfabricated channels," *PNAS* 96:5372-5377 (1999);
- Vecesey-Semjen *et al.*, "The Staphylococcal α-Toxin Pore Has a Flexible Conformation," *Biochemistry* 38 4296-4302 (1999);
- Vos de Waal *et al.* $(1977)^3$;
- Walker, B. *et al.*, "Functional Expression of the α-Hemolysin of Staphylococcus aureus in Intact Escherichia coli and in Cell Lysates," *J. Biol. Chem.* 267:10902-10909 (1992); and
- Wories *et al.*, "A novel water-soluble fluorescent probe: Synthesis, luminescence and biological properties of the sodium salt of the 4-sulfonato-3,3', 5'5-tetramethyl-2,2'-pyrromethen-1,1'-BF.sub.2 complex," *Recl. Trav. Chim. PAYSBAS* 104, 288 (1985)⁴;

Applicants have become aware of the following printed publications which may be material to the examination of this application:

- U.S. Pat. No. 4,675,285 to Clark *et al.*, provides a method for identification of clones expressing the desired protein from the cDNA libraries
- U.S. Pat. No. 5,709,998 to Kinzler *et al.*, describes APC gene, its mutations and mutations linked condition (familial adenomatous polyposis, FAP). Primers, and method for mutation detection using protein truncation test with radioactive readout is also provided.
- U.S. Pat. No. 5,861,494 to Soppet *et al.*, describes APC gene, its mutations and mutations linked condition (familial adenomatous polyposis, FAP). Primers, and method for mutation detection using protein truncation test with radioactive readout is also provided.

We have been unable to obtain this reference, but if the examiner requests a copy we will again seek to obtain it.

We have been unable to obtain this reference, but if the examiner requests a copy we will again seek to obtain it.

We have been unable to obtain this reference, but if the examiner requests a copy we will again seek to obtain it.

- U.S. Pat. No. 5,622,829 to King *et al.*, disclose the sequences of breast and ovarian cancer susceptibility genes (BRCA1). Hybridization-based methods are also described for the detection of specific mutations.
- U.S. Pat. No. 5,693,473 to Shattuck-Eidens *et al.* disclose the sequences of breast and ovarian cancer susceptibility genes (BRCA1). Hybridization-based methods are also described for the detection of specific mutations.
- U.S. Pat. No. 5,760,207 to Kinzler *et al.*, describes APC gene, its mutations and mutations linked condition (familial adenomatous polyposis, FAP). Primers, and method for mutation detection using protein truncation test with radioactive readout is also provided.
- U.S. Pat. No. 5,879,890 to Laken *et al.*, describes APC gene, its mutations and mutations linked condition (familial adenomatous polyposis, FAP). Primers, and method for mutation detection using protein truncation test with radioactive readout is also provided.
- Czworkowski *et al.*, "Fluorescence Study of the Topology of Messenger RNA Bound to the 30S Ribosomal Subunit of *Escherichia coli*," *Biochemistry* 30:4821-4830 (1991), describes the interaction of fluorescently labeled RNAs (25-36 nucleotides in length) with the fluorescently labeled 30S subunit of Escherichia coli studied by using fluorescence spectroscopic techniques.
- Hardesty *et al.*, "Ribosome function determined by fluorescence," *Biochimie* 74:391-401 (1992). This paper is a review on ribosome function.
- Picking et al., "The use of synthetic tRNA as probes for examining nascent peptides on Escherichia coli ribosomes," Biochimie 73:1101-1107 (1991), the cell free synthesis of N-acetyl or N-acyl coumarin labeled polycysteine and polyserine were carried out on Escherichia coli ribosomes using N-acyl coumarin derivatives of either Ser-tRNA or Phe-tRNA. The properties of the resulting nascent peptides were studied by fluorescence spectroscopy and compared to those of nascent polyphenylalanine chains synthesized under similar conditions.
- Picking et al., "Evidence for RNA in the Peptidyl Transferase Center of Escherichia coli Ribosomes as Indicated by Fluorescence," *Biochemistry* 31:12565-12570 (1992), The interaction of coumarin labeled (tRNA(phe)) [either the amino acid or the 5' end] with ribosomes was studied using fluorescence spectroscopy.
- Picking et al., "The Conformation of Nascent Polylysine and Polyphenylalanine Peptides on Ribosomes," J. of Biological Chemistry 266:1534-1542 (1991),

- describes the behavior of fluorescently labeled polylysine and polyphenylalanine during their in vitro synthesis on E. coli ribosomes.
- Picking et al., "Fluorescence Characterization of the Environment Encountered by Nascent Polyalanine and Polyserine as They Exit Escherichia coli Ribosomes during Translation," Biochemistry 31:2368-2375 (1992), describes a coumarin probe placed at the alpha-amino group of a synthetic elongator alanyl-tRNA or a synthetic initiator alanyl-tRNA or at the epsilon-amino group of natural lysyl-tRNA, and each was used to nonenzymatically initiate peptide synthesis.
- Picking et al., "A synthetic alanyl-initiator tRNA with initiator tRNA properties as determined by fluorescence measurements: Comparison to a synthetic alanyl-elongator tRNA," Nucleic Acids Research 19:5749-5754 (1991). A derivative of coumarin [3-(4-maleimidophenyl)-7-diethyl-amino-4-methylcoumarin] was covalently attached to the alpha amino group of alanine of the two synthetic AlatRNA species. The fluorescence spectra, quantum yield and anisotropy for the two Ala-tRNA derivatives were studied after they were bound to 70S ribosomes.
- Ma et al., "In Vitro Protein Engineering Using Synthetic tRNA^{Ala} with Different Anticodons," Biochemistry 32:7939-7945 (1993), describes the use of synthetic tRNA for in vitro protein engineering was tested in a coupled transcription/translation system prepared from Escherichia coli.
- Odom et al., "Movement of tRNA but Not the Nascent Peptide during Peptide Bond Formation on Ribosomes," Biochemistry 29:10734-10744 (1990), The interaction of fluorescently [(3-(4-maleimidophenyl)-7-diethyl-amino-4-methylcoumarin) or (5-[[2-[(iodoacetyl)amino]ethyl]amino]naphthalene-1-sulphonic acid)] labeled (Phe) tRNA at the 5'-end with ribosome as well as with nascent polypeptide was investigated using nonradiative energy transfer.
- U.S. Patent No. 5,614,386 to Metzker *et al.*, describes dyes for use in labeling DNA primers for improved DNA sequencing techniques.
- European Pat. No. 0234799A2 to Kurzchalia *et al.*, describes methods for the detection and isolation of protein utilizing the incorporation of photoaffinity reagents and biotin or other haptens into nascent peptides.
- Crowley et al., "The signal sequence moves through a ribosomal tunnel into a noncytoplasmic aqueous environment at the ER membrane early in translocation," Cell 73:1101-1115 (1993). This reference examines the logistics of nascent peptide production by utilizing ∈NBD-Lys-tRNA analogs to examine the environment of a nascent peptide chain as it moved through the ribosome and into the ER membrane.

- Karolin *et al.*, "Fluorescence and Absorption Spectroscopic Properties of Dipyrrometheneboron Difluoride (BODIPY) Derivatives in Liquids, Lipid Membranes, and Proteins," *J. Am. Chem. Soc.* 116:7801-7806 (1994) describes the use of a fluorescent dye, (BODIPY) to label a modified plasminogen activator inhibitor protein after the protein had been isolated.
- Hardesty et al., "Extension and Folding of Nascent Peptides on Ribosomes." The Translational Apparatus, Nierhaus et al. ed: New York and London; Plenum Press. p.347-358 (1993) describes peptide folding experiments where aminotRNAs were modified after aminoacylation.
- Johnson *et al.*, "Protein Synthesis and Secretion as seen by the Nascent Protein Chain," <u>The Translational Apparatus</u>, Nierhaus *et al.* ed: New York and London; Plenum Press. p. 359-370 (1993) discloses peptide synthesis experiments where Lys-tRNA was modified after aminoacylation.
- Shore *et al.*, "A Fluorescent Probe Capable of Incorporation into Nascent Polypeptide Chains," 1986 Federation Proceedings 45, 1566 Abstract, discloses the attachment of a fluorescent moiety (N⁶-fluoresceinthioacetyl) onto a tRNA aminoacylated with lysine wherein the lysine has been modified by reaction with N-bromoacetoxysuccinimide.
- Shore *et al.*, "Accessibility of AA-tRNA and Nascent Chain During Protein Synthesis," 1988 *FASEB* Journal 2, A1045 Abstract, discloses the attachment of a fluorescent moiety (N⁶-fluoresceinthioacetyl) onto a tRNA aminoacylated with lysine thereby producing N⁶-fluoresceinthioacetyl-lys-tRNA.
- Bain et al., "Site-Specific Incorporation of Non-Natural Residues into Peptides: Effect of Residue Structure on Suppression and Translation Efficiencies," *Tetrahedron* 47:2389-2400 (1991), discloses methods for the production of a series of 12 semi-synthetic acylated suppressor tRNAs.
- Johnson *et al.*, "N-Acetyllysine Transfer Ribonucleic Acid: A Biologically Active Analogue of Aminoacyl Transfer Ribonucleic Acids," *Biochemistry* 15:569-575 (1976), discloses methods for the production of N⁶-Acetyl-Lys-tRNA.
- Johnson, "Chemically Modified Aminoacyl-tRNA as a Probe of Ribosome Structure: the Synthesis and in vitro Activity of ε-N-acetyl-Lys-tRNA," 1973

 Thesis Excerpts, University of Oregon, Eugene, Oregon, discloses several methods of producing fluorescently labeled (N-methylanthranilic acid) Lys-tRNA.
- Shore, "The Use of Fluorescent-Labeled Amino Acids to Examine the Environment of Ribosome-Bound Nascent Polypeptide Chains," 1991
 Dissertation, University of Oklahoma, Norman Oklahoma, discloses the synthesis

- of fluorescent analogs of Lys-tRNA including N6-fluoresceinthioacetyl-Lys-tRNA and N⁶-6-(7-nitrobenz-2-oxa-1,3-diazol-4-yl)aminohexanoyl-Lys-tRNA.
- Kramer et al., "In Vitro engineering using synthtic tRNAs with altered anticodons including four-nucleotide anticodons," Methods Mol Biol. 77:105-16 (1998) discloses methods for the synthesis of tRNAs with altered anticodons.

Applicants have become aware of the following printed 1999 publications:

- Kramer et al., "N-terminal and C-terminal modifications affect of in vitro synthesized proteins," Int J Biochem Cell Biol. 31:231-41 (1999) discloses the effect of certain N-terminal and C-terminal modifications on protein synthesis. This publication modifies a N-acly-Met-tRNA_f with coumarin after acetylation.
- Tsalkova et al., "The effect of a hydrophobic N-terminal probe on translational pausing of chloramphenicol acetyl transferase and rhodanese," J Mol Biol. 286:71-81 (1999) discloses the effect of hydrophobic residues at the N-terminus in regards to protein synthesis. N-acetyl-S-coumarin-Met-tRNA_f was used to generate the hydrophobic residues.
- Nemoto et al., "Fluorescence labeling of the C-terminus of proteins with a puromycin analogue in cell-free translation systems," FEBS Letters. 462:43-46 (1999) discloses a method using puromycin analogues with a fluorescent moiety to label the C-terminus of nascent peptides.

This Information Disclosure Statement under 37 C.F.R. §§ 1.56 and 1.97 is not to be construed as a representation that a search has been made, that additional information material to the examination of this application does not exist, or that any one or more of these citations constitutes prior art.

Dated: August 21, 2006

Peter G. Carroll Registration No. 32,837

MEDLEN & CARROLL, LLP 101 Howard Street, Suite 350 San Francisco, California 94105 617/984-0616

FORM PTO-1449 (Modified)

AUG 2 3 2006 8

U.S. Department of Commerce Patent and Trademark Office

Attorney Docket No.: AMBER-08501

Serial No.: 10/719,523

Applicant: Kenneth J. Rothschild et al.

INFORMATION DISCLOSURE STATEMENT BY APPLICANT (Use Several Deets If Necessary)

(37 CFR § 1.98(b))

Filing Date: 11/21/03

Group Art Unit: 1636

(37 CFR § 1.9	98(b))	PADEMA		Filing Date: 11/21/03		Group Art Un	it: 1636	
	1			U.S. PATENT DOCUMENTS	Т	т	1	
Examiner Initials	Cite No.	Serial / Patent Number	Issue Date	Applicant / Patentee	Class	Subclass	Filing	g Date
	1	4,675,285	6/23/87	Clark et al.	435	6	9/1	9/84
	2	4,683,195	7/28/87	Mullis et al.	435	6	2/0	7/86
	3	4,774,339	9/27/88	Haugland <i>et al</i> .	548	405	8/1	0/87
	4	5,069,769	12/03/91	Fujimiya et al.	204	182.8	6/0	5/90
	5	5,091,328	2/25/92	Miller	437	52	11/2	1/89
	6	5,137,609	8/11/92	Manian et al.	204	180.1	1/3	1/92
•	7	5,187,288	2/16/93	Kang et al.	548 .	110	5/2:	2/91
	8	5,190,632	3/02/93	Fujimiya et al.	204	299 R	3/20)/92
-	9	5,248,782	9/28/93	Haugland et al.	548	110	12/1	8/90
	10	5,274,113	12/28/93	Kang et al.	548	405	11/0	1/91
•	11	5,433,896	7/18/95	Kang et al.	252	700	5/20)/94
	12	5,451,663	9/19/95	Kang et al.	530	367	4/08	3/93
	13	5,614,386	3/25/97	Metzker et al.	435	91.1	6/2:	3/95
	14	5,622,829	4/22/97	King et al.	435	6	4/19	9/95
	15	5,643,722	7/01/97	Rothschild et al.	435	6	5/11	/94
	16	5,451,663	9/19/95	Kang et al.	530	367	4/08	3/93
	17	5,654,150	8/05/97	King et al.	435	6	6/07	1/95
	18	5,693,473	12/02/97	Shattuck-Eidens et al.	435	6	6/07	/95
	19	5,709,998	1/20/98	Kinzler et al.	435	6	12/1	5/93
	20	5,760,207	6/02/98	Kinzler et al.	536	24.3	6/03	/96
	21	5,783,397	7/21/98	Hughes et al.	435	7.1	6/12/96	
	22	5,861,494	1/19/99	Soppet et al.	536	23.1	6/06	/95
	23	5,879,890	3/09/99	Laken et al.	435	6	1/31	/97
		<u> </u>	OREIGN PATENTS	OR PUBLISHED FOREIGN PATENT APPLICA	TIONS			
		Document	Poddiesa' D	Control (Bit is 2000)	. CI	Subclass	Trans	ation
		Document Number	Publication Date	Country / Patent Office	Class		Yes	No
	24	EP 0 234 799	02.09.87	EPO	C 12 P 21/02			
	25	WO90/05785	5/31/90	PCT				
		OTHER	DOCUMENTS (Inclu	iding Author, Title, Date, Relevant Pages, Place o	f Publication)		· <u></u>	
	26			lectric Focusing of Proteins, Walter de Gruyter, N		pp. 17-62		
	27	Antibodies: A Labor	ratory Manual (E. Har	low and D. Lane, editors, Cold Spring Harbor Lab	oratory Press, 1	988, pp. 53,72-73	3)	
	28	Bain et al., "Site-Specific Incorporation of Nonnatural Residues during In Vitro Protein Biosynthesis with Semisynthetic Aminoacyl-tRNAs," Biochemistry 30:5411-21 (1991)						
xaminer:			\/	Date Considered:				
XAMINER:	Ini	tial citation considered	. Draw line through c	itation if not in conformance and not considered.	Include conv of	this form		

FORM PTO-1449 (Modified)		U.S. Department of Commerce Patent and Trademark Office	Attorney Docket No.: AMBER-08501	Serial No.: 10/719,523	
INFORMATION DISCLOSURE STATEMENT BY APPLICANT (Use Several Sheets If Necessary)			Applicant: Kenneth J. Rothschild et al.		
(Use Several Sheets II Necessary) (37 CFR § 1.98(b))			Filing Date: 11/21/03.	Group Art Unit: 1636	
		OTHER DOCUMENTS (Including Author, Title, Da	ate, Relevant Pages, Place of Publication)		
	29	Bain et al., "Site-Specific Incorporation of Non-Natural Residue Efficiencies," <i>Tetrahedron</i> 47:2389-2400 (1991)	s into Peptides: Effect of Residue Structure	on Suppression and Translation	
	30	Bruce and Uhlenbeck, "Specific Interaction of Anticodon Loop I 3926 (1982)	Residues with Yeast Phenylalanyl-tRNA Syn	nthetase," Biochemistry 21:3921-	
	31	Crowley et al., "The signal sequence moves through a ribosomal in translocation," Cell 73:1101-1115 (1993)	tunnel into a noncytoplasmic aqueous envir	ronment at the ER membrane early	
	32	Current Protocols in Molecular Biology (F.M. Ausubel et al. ed	itors, Wiley Interscience, 1993), pp. 10-16,	10-77	
•	33	Czworkowski et al., "Fluorescence Study of the Topology of Messenger RNA Bound to the 30S Ribosomal Subunit of Escherichia coli," Biochemistry 30:4821-4830 (1991)			
	34	Da Poian, A. T., et al., "Kinetics of intracellular viral disassemble Methods 70(1), 45-58 (1998)	mbly and processing probed by Bodipy fluorescence dequenching," J Virol		
	35	Doty et al., "Strand Separation and Specific Recombination in D USA 46:461-476 (1960)	ion in Deoxyribonucleic Acids: Physical Chemicals Studies," Proc. Natl. Acad. Sci.		
•	36	DiCesare et al., "A High-Sensitivity Electrochemiluminescence- BioTechniques 15:152-59 (1993)	Based Detection System for Automated PCI	R Product Quantitation,"	
	37	Felgner et al., "Lipofection: A highly efficient, lipid-mediated Di	NA-transfection procedure," Proc. Natl. Acc	ad. Sci. USA 84:7413-17 (1987)	
	38	Happ et al., "New Approach to the Synthesis of 2'(3')-O-Aminoa	cyl Oligoribonucleotides," J. Org. Chem. 52	2:5387-91 (1987)	
	39	Hardesty et al., "Extension and Folding of Nascent Peptides on Ribosomes." The Translational Apparatus, Nierhaus et al. ed: New York and London; Plenum Press. p. 347-358 (1993)			
	40	Hardesty et al., "Ribosome function determined by fluorescence,	" Biochimie 74:391-401 (1992)		
	41	Heckler et al., "Preparation of 2'(3')-O-Acyl-pCpA Derivatives a: Tetrahedron 40:87-94 (1984)	s Substrates for T4 RNA Ligase-Mediated "(Chemical Aminoacylation","	
	42	Heckler et al., "T4 RNA Ligase Mediated Preparation of Novel "Chemically Misacylated" tRNA Phes," Biochemistry 23:1468-73 (1984)			
	43	Hemmila, I.A., Chemical Analysis "Applications of Fluorescence in Immunoassays", (Wiley&Sons 1991) pp.138-159			
	44	Hudson, "Methodological Implications of Simultaneous Solid-Phases Peptide Synthesis. 1. Comparison of Different Coupling Procedures," J. Org. Chem. 53:617-624 (1988)			
	45	Johnson et al., "Protein Synthesis and Secretion as seen by the N York and London; Plenum Press. p. 359-370 (1993)	ascent Protein Chain," The Translational Ap	paratus, Nierhaus et al. ed: New	
	46	Johnson et al., "N-Acetyllysine Transfer Ribonucleic Acid: A Bio Biochemistry 15:569-575 (1976)	ologically Active Analogue of Aminoacyl Tr	ransfer Ribonucleic Acids,"	
	47	Johnson, "Chemically Modified Aminoacyl-tRNA as a Probe of ItRNA," 1973 Thesis Excerpts, University of Oregon, Eugene, Or	Ribosome Structure: the Synthesis and in vit regon	tro Activity of ε-N-acetyl-Lys-	
	48	Karolin et al., "Fluorescence and Absorption Spectroscopic Prop. Lipid Membranes, and Proteins," J. Am. Chem. Soc. 116:7801-78	ertes of Dipyrrometheneboron Difluoride (B 806 (1994)	ODIPY) Derivatives in Liquids,	
	49	Keller, R. C., et al., "Characterization of the Resonance Energy T Lipid Research," Biochem Biophys Res Commun 207(2), 508-14	Transfer Couple Coumarin-Bodipy and its Po (1995)	ossible Applications in Protein-	
	50	Kim, D., and Choi, C., "A Semicontinuous Prokaryotic Coupled Transcription/Translation System Using a Dialysis Membrane," Biotechnol Prog 12, 645-649 (1996)			
	51	Kopp et al., "Chemical Amplification: Continuous Flow PCR on a Chip," Science 280:1046 (1998)			
	52	Kozak, "Point Mutations Define a Sequence Flanking the AUG Initiator Codon that Modulates Translation by Eukaryotic Ribosomes," Cell 44:283-292 (1986)			
	53	Kramer et al., "In Vitro engineering using synthtic tRNAs with altered anticodons including four-nucleotide anticodons," Methods Mol Biol. 77:105-16 (1998)			
	54	Kramer et al., "N-terminal and C-terminal modifications affect of	f in vitro synthesized proteins," Int J Biocher	m Cell Biol. 31:231-41 (1999)	
	55	Krieg et al., "Photocrosslinking of the signal sequence of nascent particle," <i>Proc. Natl. Acad. Sci. USA</i> 83:8604-08 (1986)	preprolactin to the 54-kilodalton polypeptid	le of the signal recognition	
	56	Kudlicki, W.et al., "Chaperone-dependent Folding and Activation	n of Ribosome-bound Nascent Rhodanse,"	Mol Biol 244(3), 319-31 (1994)	
	57	Laemmli, U. K., "Cleavage of Structural Proteins during the Asse	embly of the Head of Bacteriophage T4," No	ature 227:680-685 (1970)	
Examiner:			Date Considered:		
EXAMINER:	Initi	ial citation considered. Draw line through citation if not in conform	nance and not considered. Include copy of the	his form	

FORM PTO-144 (Modified)	FORM PTO-1449 Modified) U.S. Department of Commerce Patent and Trademark Office		Attorney Docket No.: AMBER-08501	Serial No.: 10/719,523	
INFORMATION DISCLOSURE STATEMENT BY APPLICANT (Use Several Sheets If Necessary)			Applicant: Kenneth J. Rothschild et al.		
(37 CFR § 1.98(b))			Filing Date: 11/21/03	Group Art Unit: 1636	
		OTHER DOCUMENTS (Including Author, Title, D.			
	Ma et al., "In Vitro Protein Engineering Using Synthetic tRNA ha with Different Anticodons," Biochemistry 32:7939-7945 (1993)				
	Marmur and Lane, "Strand Separation and Specific Recombination in Deoxyribonucleic Acids: Biological Studies," <i>Proc. Natl. Acad. S</i> 46:453-461 (1960) Molecular Cell Biology (J. Darnell et al. editors, Scientific American Books, N.Y., N.Y. 1991) pp. 119-132				
•	61	402.43-40 (1999)			
	Neu and Heppel, "Nucleotide Sequence Analysis of Polyribonucleotides by Means of Periodate Oxidation Followed by Cleavage with an Amine," J. Biol. Chem. 239:2927-34 (1964)				
	Noren et al., "A General Method for Site-Specific Incorporation of Unnatural Amino Acids into Proteins," Science 244:182-188 (1989)				
	Odom, O. W, et al., "In vitro engineering using acyl-derivatized tRNA," In <u>Protein synthesis: Methods and Protocols</u> , PP.93-103, (Humpress, Totowa, NJ.)				
	Odom et al., "Movement of tRNA but Not the Nascent Peptide during Peptide Bond Formation on Ribosomes," Biochemistry 29:10734-10 (1990)				
	66 Patchornik et al., "Photosensitive Protecting Groups," J. Am. Chem. Soc. 92:6333-35 (1970)				
	Pavlopoulos, et al., "Laser action from a tetramethylpyrromethene-BF.sub.2 complex," APP. OPTICS 27:4998-4999 (1988)				
	Picking et al., "The use of synthetic tRNA as probes for examing nascent peptides on Escherichia coli ribosomes," Biochimie 73:1101-110 (1991)				
	69	Picking et al., "Evidence for RNA in the Peptidyl Transferase Co 31:12565-12570 (1992)	enter of Escherichia coli Ribosomes as Indic	ated by Fluorescence," Biochemist.	
	70	Picking <i>et al.</i> , "The Conformation of Nascent Polylysine and Pol 1542 (1991)	yphenylalanine Peptides on Ribosomes," J.	of Biological Chemistry 266:1534	
	71	Picking et al., "Fluorescence Characterization of the Environme coli Ribosomes during Translation," Biochemistry 31:2368-2375	nt Encountered by Nascent Polyalanine and (1992)	Polyserine as They Exit Escherichi	
	72	Picking et al., "A synthetic alanyl-initiator tRNA with initiator tRNA properties as determined by fluorescence measurements: Comparison to synthetic alanyl-elongator tRNA," Nucleic Acids Research 19:5749-5754 (1991)			
	73	Pillai, "Photoremovable Protecting Groups in Organic Synthesis," Synthesis 1-26 (1980)			
	Powell et al., "Molecular Diagnosis of Familial Adenomatous Polyposis," N. Engl. J. Med. 329:1982-87 (1993)				
	75	Pratt, "Coupled Transcription-Translation in Prokaryotic Cell-Freeditors, p. 179-209, IRL Press, Oxford, 1984)	ee System," (Transcription and Translation	, B.D. Hames and S.J. Higgins,	
	76	Promega Technical Bulletin No. 182; tRNA nscendTM: Non-radioact	tive Translation Detection System, Sept. 199	93	
	77	Reis, R. C., et al., "A novel methodology for the investigation of (1998)	intracellular proteolytic processing in intet	cells," Eur J Cell Biol 75(2), 192-7	
	Rowan and Bodmer, "Introduction of a myc Reporter Taq to Improve the Quality of Mutation Detection Using the Protein Truncation Test," Human Mutation 9:172-176 (1997)				
	Sampson and Uhlenbeck, "Biochemical and physical characterization of an unmodified yeast phenylalanine transfer RNA transcribed in vitre Proc. Natl. Acad. Sci. USA 85:1033-37 (1988)				
	80	Seong and RajBhandary, "Escherichia coli formylmethione tRNA affect initiation of protein synthesis and conformation of anticod	A: Mutations in ^{GGG} sequence conserved in a on loop," <i>Proc. Natl. Acad. Sci. USA</i> 84:334	nticodon stem of initiator tRNAs I-338 (1987)	
	81	Shore et al., "A Fluorescent Probe Capable of Incorporation into	Nascent Polypeptide Chains," 1986 Federa	tion Proceedings 45, 1566 Abstrac	
	82	Shore et al., "Accessibility of AA-tRNA and Nascent Chain Duri	ing Protein Synthesis," 1988 FASEB Journal	2, A1045 Abstract	
	83	Shore, "The Use of Fluorescent-Labeled Amino Acids to Examin Dissertation, University of Oklahoma, Norman Oklahoma	e the Environment of Ribosome-Bound Nas	cent Polypeptide Chains," 1991	
	84	Spirin et al., "A Continous Cell-Free Translation System Capable			
	85	Stephens, "High-Resolution Preparative SDS-Polyacrylamide Ge Concentration of Protein Bands," Anal. Biochem. 65:369-79 (197	l Electrophoresis: Fluorescent Visualization 75)	and Electrophoretic Elution-	
	86	Treibs & Kreuzer, "Difluorboryl-komplexe von di- und tripyrryln	nethenen," Liebigs Ann. Chem. 718:208-223	(1968)	
			Date Considered:		
Examiner:					

FORM PTO-1449 (Modified)	U.S. Department of Commerce Patent and Trademark Office	Attorney Docket No.: AMBER-08501	Serial No.: 10/719,523		
' '	TION DISCLOSURE STATEMENT BY APPLICANT (Use Several Sheets If Necessary)	Applicant: Kenneth J. Rothschild et al.			
(37 CFR § 1.98(b))	(Use Several Sheets if Necessary)	Filing Date: 11/21/03	Group Art Unit: 1636		
	OTHER DOCUMENTS (Including Author, Title, I	Date, Relevant Pages, Place of Publication)			
87	Tsalkova et al., "The effect of a hydrophobic N-terminal probe Mol Biol. 286:71-81 (1999)	on translational pausing of chloramphenicol	acetyl transferase and rhodanese," J		
88	Turcatti et al., "Probing the Structure and Function of the Tachykinin Neurokinin-2 Receptor through Biosynthetic Incorporation of Fluorescent Amino Acids at Specific Sites," J Biol Chem 271(33), 19991-8 (1996)				
89	Varshney U, RajBhandary UL, "Initiation of protein synthesis from a termination codon," Proc Natl Acad Sci U S A 87(4):1586-90 (1990)				
90	Varshney et al., "Direct Analysis of Aminoacylation Levels of	Varshney et al., "Direct Analysis of Aminoacylation Levels of tRNAa in Vivo," J. Biol. Chem. 266: 24712-24718 (1991)			
91	Vecesey-Semjen et al., "The Staphylococcal α-Toxin Pore Has a Flexible Conformation," Biochemistry 38 4296-4302 (1999)				
92	Walker, B. et al., "Functional Expression of the α-Hemolysin of Staphylococcus aureus in Intact Escherichia coli and in Cell Lysates," J. Biol. Chem. 267:10902-10909 (1992)				
93	Yao S et al., "SDS capillary gel electrophoresis of proteins in microfabricated channels," PNAS 96:5372-5377 (1999)				
94					
95					
96					
. 97					
98					
99					
100					
101					
102					
103			· · · · · · · · · · · · · · · · · · ·		
104					
105					
106					
107			····		
108					
109					
110					
111					
Examiner:		Date Considered:			
EXAMINER:	EXAMINER: Initial citation considered. Draw line through citation if not in conformance and not considered. Include copy of this form				

This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record.

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

BLACK BORDERS

IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

FADED TEXT OR DRAWING

BLURRED OR ILLEGIBLE TEXT OR DRAWING

SKEWED/SLANTED IMAGES

COLOR OR BLACK AND WHITE PHOTOGRAPHS

GRAY SCALE DOCUMENTS

LINES OR MARKS ON ORIGINAL DOCUMENT

REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

IMAGES ARE BEST AVAILABLE COPY.

OTHER:

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.